



PROPTOSIS REPLACEMENT



BEFORE




AFTER


This dog experienced a traumatic proptosis after being bitten by a larger dog. The globe was surgically replaced into the socket.



CORNEAL DERMOID REMOVAL



BEFORE




AFTER


This patient had a corneal dermoid, which is a congenital malformation resulting in skin and hair growing from the ocular surface. Surgical resection of the abnormal area yielded excellent clarity and comfort.



CORNEAL DERMOID REMOVAL



BEFORE




AFTER

This patient had a corneal dermoid, which is a congenital malformation resulting in skin and hair growing from the ocular surface. Surgical resection of the abnormal area yielded excellent clarity and comfort.

KERATITIS TREATMENT



BEFORE




AFTER


This cat developed immune-mediated keratitis, which responded favorably to medical therapy.



KERATITIS TREATMENT



BEFORE



AFTER

This cat developed immune-mediated keratitis, which responded favorably to medical therapy.



EYELID RECONSTRUCTION



BEFORE



AFTER

This kitten was born without upper eyelids, resulting in hair contacting the corneal surface and pain. Tissue grafting was performed to create a new functioning upper eyelid for each eye.



EYELID RECONSTRUCTION



BEFORE



AFTER

This kitten was born without upper eyelids, resulting in hair contacting the corneal surface and pain. Tissue grafting was performed to create a new functioning upper eyelid for each eye.

